

Medicines & Healthcare products Regulatory Agency

> MHRA 10 South Colonnade Canary Wharf London E14 4PU United Kingdom

gov.uk/mhra

## **Decision Cover Letter**

### Decision of the licensing authority to:

accept change(s) to the agreed paediatric investigation plan (MHRA-100623-PIP01-22-M01) and to the deferral

MHRA-100623-PIP01-22-M02

### Scope of the Application

### **Active Substance(s)**

NIRAPARIB TOSYLATE MONOHYDRATE

### **Condition(s)**

Treatment of all conditions included in the category of malignant neoplasms (except haematopoietic and lymphoid malignancies)

### **Pharmaceutical Form(s)**

Capsule, hard; Tablet; Tablet for oral suspension

### **Route**(s) of Administration

ORAL USE

### Name / Corporate name of the PIP applicant

GlaxoSmithKline UK Limited

### **Basis for the Decision**

Pursuant to the Human Medicines Regulations 2012, GlaxoSmithKline UK Limited submitted to the licensing authority on 21/03/2024 12:39 GMT an application for a

The procedure started on 04/06/2024 10:50 BST

1. The licensing authority, having assessed the application in accordance with the Human Medicines Regulations 2012, decides, as set out in the appended summary report:

to accept change(s) to the agreed paediatric investigation plan and to the deferral

2. The measures and timelines of the paediatric investigation plan are set out in the Annex I.

This decision is forwarded to the applicant, together with its annex and appendix.



Medicines & Healthcare products Regulatory Agency

> MHRA 10 South Colonnade Canary Wharf London E14 4PU United Kingdom

gov.uk/mhra

# **Final Decision Letter**

MHRA-100623-PIP01-22-M02

Of 03/09/2024 06:50 BST

On the adopted decision for NIRAPARIB TOSYLATE MONOHYDRATE (MHRA-100623-PIP01-22-M02) in accordance with the Human Medicines Regulations 2012.

The licensing authority, in accordance with the Human Medicines Regulations 2012, has adopted this decision:

Agreement on modification of a paediatric investigation plan (including modification of a waiver or deferral included in that paediatric investigation plan)

This decision applies to a for NIRAPARIB TOSYLATE MONOHYDRATE, Capsule, hard; Tablet; Tablet for oral suspension, ORAL USE.

This decision is addressed to GlaxoSmithKline UK Limited, 980 Great West Road, Brentford, UNITED KINGDOM, TW8 9GS

# ANNEX I

1. Waiver

### **1.1 Condition:**

Not applicable

### 2. Paediatric Investigation Plan:

### 2.1 Condition(s):

Treatment of all conditions included in the category of malignant neoplasms (except haematopoietic and lymphoid malignancies).

### **2.2 Indication(s) targeted by the PIP:**

Treatment of paediatric patients from birth to less than 18 years old with neuroblastoma and/or osteosarcoma

### **2.3** Subset(s) of the paediatric population concerned by the paediatric development:

The paediatric population from birth to less than 18 years of age

### **2.4 Pharmaceutical Form(s):**

Film coated tablet Capsule, hard Table for oral suspension

### 2.5 Studies:

Quality Measures 1 Study 1 Development of age- appropriate oral liquid dosage form.   Non-Clinical Studies 0 Not applicable.   Clinical Studies 3 Study 2 Open-label, multiple dose, two part trial to evaluate pharmacokinetics, safety, activity and acceptability of niraparib when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18 years of age with relapsed/ refractory osteosarcoma	Study Type	Number of Studies	Study Description
Non-Clinical Studies0Not applicable.Clinical Studies3Study 2 Open-label, multiple dose, two part trial to evaluate pharmacokinetics, safety, activity and acceptability of niraparib when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age to less than 18 years	Quality Measures	1	Study 1 Development of age-
Clinical Studies3Study 2 Open-label, multiple dose, two part trial to evaluate pharmacokinetics, safety, activity and acceptability of niraparib when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age to less than 18 years of age with recurrent / refractory neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib 			
dose, two part trial to evaluate pharmacokinetics, safety, activity and acceptability of niraparib when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18	Non-Clinical Studies	-	
pharmacokinetics, safety, activity and acceptability of niraparib when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18	Clinical Studies	3	Study 2 Open-label, multiple
and acceptability of niraparib when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
when given in combination with dostarlimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
dostarĪimab in children from 6 months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			and acceptability of niraparib
months of age to less than 18 years of age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
age with recurrent / refractory solid tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
tumours, excluding central nervous system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
system (CNS) tumours in part 1a and 1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
1b and with recurrent / refractory osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
osteosarcoma and recurrent / refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
refractory neuroblastoma in Part 2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
2. Study 3 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
of niraparib in combination with dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
dostarlimab against current standard of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
of care in children from 6 months of age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
age to less than 18 years of age with relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
relapsed/ refractory osteosarcoma and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
and/or neuroblastoma. Study 4 Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
Open label, randomised controlled, active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
active comparator trial to evaluate efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
efficacy and safety of niraparib in combination with dostarlimab against current standard of care in children from birth to less than 18			
in combination with dostarlimab against current standard of care in children from birth to less than 18			
against current standard of care in children from birth to less than 18			
children from birth to less than 18			
years of age with newly diagnosed			years of age with newly diagnosed

	1	high risk osteosarcoma and/or Stage 4 neuroblastoma.
Extrapolation, Modeling &	1	Study 5 Modelling and simulation
Simulation Studies		study, to evaluate the use of niraparib
		and dostarlimab in the proposed
		paediatric indications in children
		from birth to less than 18 years of
		5
		age.
Other Studies	0	Not applicable.
Other Measures	0	Not applicable.

# 3. Follow-up, completion and deferral of a PIP:

Concerns on potential long term safety and efficacy issues in relation to paediatric use:	Yes
Date of completion of the paediatric investigation plan:	30/06/2040
Deferral of one or more studies contained in the paediatric investigation plan:	Yes